



Johnson P, Federico M, Kirkwood A, et al. Adapted treatment guided by interim PET-CT scan in advanced Hodgkin's Lymphoma. *New Engl J Med*. 2016;374: 2419-2429.

Engert A, Haverkamp H, Kobe C, et al. Reduced-intensity chemotherapy and PET-guided radiotherapy in patients with advanced stage Hodgkin's lymphoma (HD15 trial): a randomised, open-label, phase 3 noninferiority trial. *Lancet* 2012;379:1791-1799.

Casasnovas RO, Bouabdallah R, Brice P, et al. PET-adapted treatment for newly diagnosed advanced Hodgkin lymphoma (AHL2011): a randomised, multicentre, noninferiority, phase 3 study. *Lancet Oncol* 2019;20:202-215.

Casasnovas RO, Bouabdallah R, Brice P, et al. Positron Emission Tomography-Driven Strategy in Advanced Hodgkin Lymphoma: Prolonged Follow-Up of the AHL2011 Phase III Lymphoma Study Association Study. *J Clin Oncol*. 2022 Apr 1;40(10):1091-1101. PMID: 34990281.

Borchmann P, Goergen H, Kobe C et al. PET-guided treatment in patients with advanced-stage Hodgkin's lymphoma (HD18): final results of an open-label, international, randomised phase 3 trial by the German Hodgkin Study Group. *Lancet*. 2017 Dec 23;390(10114):2790-2802. doi: 10.1016/S0140-6736(17)32134-7. Epub 2017 Oct 20. PMID: 29061295.

Kreissl S, Goergen H, Buehnen I et al. German Hodgkin Study Group. PET-guided eBEACOPP treatment of advanced-stage Hodgkin lymphoma (HD18): follow-up analysis of an international, open-label, randomised, phase 3 trial. *Lancet Haematol*. 2021 Jun;8(6):e398-e409.

von Tresckow B, Plütschow A, Fuchs M, et al. Dose-intensification in early unfavorable Hodgkin's lymphoma: final analysis of the German Hodgkin Study Group HD14 trial. *J Clin Oncol* 2012;30:907-913.

Straus DJ, Długosz-Danecka M, Connors JM, et al. Brentuximab vedotin with chemotherapy for stage III or IV classical Hodgkin lymphoma (ECHELON-1): 5-year update of an international, open-label, randomised, phase 3 trial. *Lancet Haematol* 2021;8:e410-e421.

Savage KJ, Skinnider B, Al-Mansour M, et al. Treating limited-stage nodular lymphocyte predominant Hodgkin lymphoma similarly to classical Hodgkin lymphoma with ABVD may improve outcome. *Blood*. 2011 Oct 27;118(17):4585-90. doi: 10.1182/blood-2011-07-365932. Epub 2011 Aug 26. PMID: 21873543.

Canellos GP, Mauch P. What is the appropriate systemic chemotherapy for lymphocyte-predominant Hodgkin's lymphoma? *J Clin Oncol*. 2010 Jan 1;28(1):e8. doi: 10.1200/JCO.2009.26.3293. Epub 2009 Nov 23. PMID: 19933898.

Fanale MA, Cheah CY, Rich A, et al. Encouraging activity for R-CHOP in advanced stage nodular lymphocyte-predominant Hodgkin lymphoma. *Blood*. 2017 Jul 27;130(4):472-477. doi: 10.1182/blood-2017-02-766121. Epub 2017 May 18. PMID: 28522441; PMCID: PMC5578726.

Shankar A, Hall GW, Gorde-Grosjean S, et al. Treatment outcome after low intensity chemotherapy [CVP] in children and adolescents with early stage nodular lymphocyte predominant Hodgkin's lymphoma - an AngloFrench collaborative report. *Eur J Cancer*. 2012 Jul;48(11):1700-6. doi: 10.1016/j.ejca.2011.10.018. Epub 2011 Nov 15. PMID: 22093944.

Advani RH, Horning SJ, Hoppe RT, Daadi S, Allen J, Natkunam Y, Bartlett NL. Mature results of a phase II study of rituximab therapy for nodular lymphocyte-predominant Hodgkin lymphoma. *J Clin Oncol*. 2014 Mar 20;32(9):912-8. doi: 10.1200/JCO.2013.53.2069. Epub 2014 Feb 10. PMID: 24516013.

Note: Pathways are independent of specific OptumCare/health plan medical policy coverage criteria. OptumCare/health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars or alternate formulations (along with the reference products) are considered on pathway unless otherwise specified by OptumCare/health plan formularies, medical policies, or preferred product rules.

Advani RH, Hoppe RT. How I treat nodular lymphocyte predominant Hodgkin lymphoma. *Blood*. 2013 Dec 19;122(26):4182-8. doi: 10.1182/blood-2013-07-453241. Epub 2013 Nov 8. PMID: 24215035.

Eichenauer DA, Fuchs M, Plütschow A, et al. Phase 2 study of rituximab in newly diagnosed stage IA nodular lymphocyte-predominant Hodgkin lymphoma: a report from the German Hodgkin Study Group. *Blood*. 2011 Oct 20;118(16):4363-5. doi: 10.1182/blood-2011-06-361055. Epub 2011 Aug 9. PMID:21828141.

Eichenauer DA, Plütschow A, Fuchs M, et al. Long-Term Course of Patients With Stage IA Nodular Lymphocyte-Predominant Hodgkin Lymphoma: A Report From the German Hodgkin Study Group. *J Clin Oncol*. 2015 Sep 10;33(26):2857-62. doi: 10.1200/JCO.2014.60.4363. Epub 2015 Aug 3. PMID: 26240235.

Schulz H, Rehwald U, Morschhauser F, et al. Rituximab in relapsed lymphocyte-predominant Hodgkin lymphoma: long-term results of a phase 2 trial by the German Hodgkin Lymphoma Study Group (GHSG). *Blood*. 2008 Jan 1;111(1):109-11. doi: 10.1182/blood-2007-03-078725. Epub 2007 Oct 15. PMID: 17938252.

Engert A, Plütschow A, Eich HT, et al. Reduced treatment intensity in patients with early-stage Hodgkin's lymphoma. *N Engl J Med*. 2010 Aug 12;363(7):640-52. doi: 10.1056/NEJMoa1000067. PMID: 20818855.

Stamatoullas A, Brice P, Bouabdallah R, et al. Outcome of patients older than 60 years with classical Hodgkin lymphoma treated with front line ABVD chemotherapy: frequent pulmonary events suggest limiting the use of bleomycin in the elderly. *Br J Haematol*. 2015 Jul;170(2):179-84. doi: 10.1111/bjh.13419. Epub 2015 Apr 19. PMID: 25891777.

Behringer K, Goergen H, Hitz F, et al; German Hodgkin Study Group; Swiss Group for Clinical Cancer Research; Arbeitsgemeinschaft Medikamentöse Tumortherapie. Omission of dacarbazine or bleomycin, or both, from the ABVD regimen in treatment of early-stage favourable Hodgkin's lymphoma (GHSG HD13): an open-label, randomised, non-inferiority trial. *Lancet*. 2015 Apr 11;385(9976):1418-27. doi: 10.1016/S0140-6736(14)61469-0. Epub 2014 Dec 22. Erratum in: *Lancet*. 2015 Apr 11;385(9976):1396. PMID: 25539730.

Kolstad A, Nome O, Delabie J, et al. Standard CHOP-21 as first line therapy for elderly patients with Hodgkin's lymphoma. *Leuk Lymphoma*. 2007 Mar;48(3):570-6. doi: 10.1080/10428190601126610. PMID: 17454601.

Evens AM, Advani RH, Helenowski IB, et al. Multicenter Phase II Study of Sequential Brentuximab Vedotin and Doxorubicin, Vinblastine, and Dacarbazine Chemotherapy for Older Patients With Untreated Classical Hodgkin Lymphoma. *J Clin Oncol*. 2018 Oct 20;36(30):3015-3022. doi: 10.1200/JCO.2018.79.0139. Epub 2018 Sep 4. PMID: 30179569.

Friedberg JW, Forero-Torres A, Bordoni RE, et al. Frontline brentuximab vedotin in combination with dacarbazine or bendamustine in patients aged  $\geq 60$  years with HL. *Blood*. 2017 Dec 28;130(26):2829-2837. doi: 10.1182/blood-2017-06-787200. Epub 2017 Oct 16. PMID: 29038340.

Younes A, Gopal AK, Smith SE, et al. Results of a pivotal phase II study of brentuximab vedotin for patients with relapsed or refractory Hodgkin's lymphoma. *J Clin Oncol* 2012;30:2183-2189.

O' Connor OA, Lue JK, Sawas A, et al. Brentuximab vedotin plus bendamustine in relapsed or refractory Hodgkin's lymphoma: an international, multicentre, single-arm, phase 1-2 trial. *Lancet Oncol* 2018;19:257-266.

Note: Pathways are independent of specific OptumCare/health plan medical policy coverage criteria. OptumCare/health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars or alternate formulations (along with the reference products) are considered on pathway unless otherwise specified by OptumCare/health plan formularies, medical policies, or preferred product rules.

Herrera AF, Moskowitz AJ, Bartlett NL, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with relapsed or refractory Hodgkin lymphoma. *Blood* 2018;131:1183-1194.

Josting A, Rudolph C, Reiser M, et al. Time-intensified dexamethasone/cisplatin/ cytarabine: an effective salvage therapy with low toxicity in patients with relapsed and refractory Hodgkin's disease. *Ann Oncol* 2002;13:1628-1635.

Aparicio J, Segura A, Garcera S, et al. ESHAP is an active regimen for relapsing Hodgkin's disease. *Ann Oncol* 1999;10:593-595.

Labrador J, Cabrero-Calvo M, Perez-Lopez E, et al. ESHAP as salvage therapy for relapsed or refractory Hodgkin's lymphoma. *Ann Hematol* 2014;93:1745-1753.

Fernández de Larrea C, Martínez C, et al. Salvage chemotherapy with alternating MINE-ESHAP regimen in relapsed or refractory Hodgkin's lymphoma followed by autologous stem cell transplantation. *Ann Oncol* 2010;21:1211-1216.

Santoro A, Mazza R, Pulsoni A, et al. Bendamustine in combination with gemcitabine and vinorelbine is an effective regimen as induction chemotherapy before autologous stem-cell transplantation for relapsed or refractory Hodgkin lymphoma: final results of a multicenter phase II study. *J Clin Oncol* 2016;34:3293-3299.

Bartlett N, Niedzwiecki D, Johnson J, et al. Gemcitabine, vinorelbine, and pegylated liposomal doxorubicin (GVD), a salvage regimen in relapsed Hodgkin's lymphoma: CALGB 59804. *Ann Oncol* 2007;18:1071-1079.

Moskowitz AJ, Shah G, Schöder H, et al. Phase II trial of pembrolizumab plus gemcitabine, vinorelbine, and liposomal doxorubicin as second-line therapy for relapsed or refractory classical Hodgkin lymphoma. *J Clin Oncol* 2021;39:3109-3117.

Moskowitz CH, Nimer SD, Zelenetz AD, et al. A 2-step comprehensive high-dose chemoradiotherapy second-line program for relapsed and refractory Hodgkin disease: analysis by intent to treat and development of a prognostic model. *Blood* 2001;97:616-623.

Hertzberg MS, Crombie C, Benson W, et al. Outpatient fractionated ifosfamide, carboplatin and etoposide as salvage therapy in relapsed and refractory non-Hodgkin's and Hodgkin's lymphoma. *Ann Oncol* 2006;17 Suppl 4:iv25-30.

Abali H, Urün Y, Oksüzoğlu B, Budakoğlu B, et al. Comparison of ICE (ifosfamidecarboplatin-etoposide) versus DHAP (cytosine arabinoside-cisplatin-dexamethasone) as salvage chemotherapy in patients with relapsed or refractory lymphoma. *Cancer Invest* 2008;26:401-406.Santoro 2007

Chen R, Zinzani PL, Fanale MA, et al. Phase II study of the efficacy and safety of pembrolizumab for relapsed/refractory classic Hodgkin lymphoma. *J Clin Oncol* 2017;35:2125-2132.

Kuruvilla J, Ramchandren R, Santoro A, et al. Pembrolizumab versus brentuximab vedotin in relapsed or refractory classical Hodgkin lymphoma (KEYNOTE-204): an interim analysis of a multicentre, randomised, open-label, phase 3 study. *Lancet Oncol* 2021;22:512-524.

Note: Pathways are independent of specific OptumCare/health plan medical policy coverage criteria. OptumCare/health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars or alternate formulations (along with the reference products) are considered on pathway unless otherwise specified by OptumCare/health plan formularies, medical policies, or preferred product rules.